

LYNPARZA is the only PARPi FDA-approved for monotherapy and combination therapy in advanced ovarian cancer¹⁻³



Treatment setting	Prior lines and response to chemotherapy	Required biomarker status per indication*	FDA-approved PARP inhibitor
First-Line Maintenance Treatment	• ≥1 prior line • Complete response (CR) or partial response (PR) to platinum-based chemotherapy	HRD-positive status defined by: • a <i>BRCA</i> mutation, and/or • genomic instability	LYNPARZA® (olaparib) in combination with bevacizumab
		Germline or somatic <i>BRCA</i> mutation	LYNPARZA® (olaparib) monotherapy
		HRD-positive status defined by: • a <i>BRCA</i> mutation, and/or • genomic instability	Niraparib
Maintenance Treatment for Recurrence	• ≥2 prior lines • CR or PR to platinum-based chemotherapy	Germline <i>BRCA</i> mutation	Niraparib
		Germline or somatic <i>BRCA</i> mutation	LYNPARZA® (olaparib) monotherapy
			Rucaparib

No safety and/or efficacy comparisons should be made between these agents as no head-to-head clinical trials have been conducted.

*As detected by the appropriate FDA-approved companion diagnostic. Refers to biomarkers listed within indications.

LYNPARZA is a poly (ADP-ribose) polymerase (PARP) inhibitor indicated:

- for the maintenance treatment of adult patients with deleterious or suspected deleterious germline or somatic *BRCA*-mutated (g*BRCA*m or s*BRCA*m) advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in complete or partial response to first-line platinum-based chemotherapy. Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA
- in combination with bevacizumab for the maintenance treatment of adult patients with advanced epithelial ovarian, fallopian tube or primary peritoneal cancer who are in complete or partial response to first-line platinum-based chemotherapy and whose cancer is associated with homologous recombination deficiency (HRD)-positive status defined by either a deleterious or suspected deleterious *BRCA* mutation, and/or genomic instability. Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA
- for the maintenance treatment of adult patients with deleterious or suspected deleterious germline or somatic *BRCA*-mutated (g*BRCA*m or s*BRCA*m) recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer, who are in complete or partial response to platinum-based chemotherapy. Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA

See Prescribing Information for LYNPARZA, Zejula, and Rubraca for full indications, including which companion diagnostics to use with each treatment.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS

There are no contraindications for LYNPARZA.

WARNINGS AND PRECAUTIONS

Myelodysplastic Syndrome/Acute Myeloid Leukemia (MDS/AML): Occurred in approximately 1.2% of patients (26/2219) with various *BRCA*m, g*BRCA*m, HRR gene-mutated or HRD-positive cancers who received LYNPARZA in clinical studies as a single agent or as part of a combination regimen, consistent with the approved indications, and the majority of events had a fatal outcome. The median duration of therapy in patients who developed MDS/AML was approximately 2 years (range: <6 months to >4 years). All of these patients had previous chemotherapy with platinum agents and/or other DNA-damaging agents, including radiotherapy.

In SOLO-1, patients with newly diagnosed advanced *BRCA*m ovarian cancer, the incidence of MDS/AML was 1.9% (5/260) in patients who received LYNPARZA and 0.8% (1/130) in patients who received placebo based on

Please see additional Important Safety Information on the next page and complete Prescribing Information, including Medication Guide.

*BRCA*m=*BRCA*-mutated; g*BRCA*m=germline *BRCA*-mutated; HRD=homologous recombination deficiency; HRR=homologous recombination repair; PARPi=poly (ADP-ribose) polymerase inhibitor; s*BRCA*m=somatic *BRCA*-mutated.

Niraparib (Zejula®) is indicated²:

- for the maintenance treatment of adult patients with advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in a complete or partial response to first-line platinum-based chemotherapy
- for the maintenance treatment of adult patients with deleterious or suspected deleterious germline *BRCA*-mutated recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in a complete or partial response to platinum-based chemotherapy. Select patients for therapy based on an FDA-approved companion diagnostic for ZEJULA

Rucaparib (Rubraca®) is indicated³:

- for the maintenance treatment of adult patients with a deleterious *BRCA* mutation (germline and/or somatic)-associated recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in a complete or partial response to platinum-based chemotherapy. Select patients for therapy based on an FDA-approved companion diagnostic for RUBRACA

an updated analysis. In PAOLA-1, of patients with newly diagnosed advanced ovarian cancer with HRD-positive status, the incidence of MDS/AML was 1.6% (4/255) in patients who received LYNPARZA and 2.3% (3/131) in the control arm.

In SOLO-2, patients with *BRCA*m platinum-sensitive relapsed ovarian cancer, the incidence of MDS/AML was 8% (15/195) in patients who received LYNPARZA and 4% (4/99) in patients who received placebo. The duration of LYNPARZA treatment prior to the diagnosis of MDS/AML ranged from 0.6 years to 4.5 years.

Do not start LYNPARZA until patients have recovered from hematological toxicity caused by previous chemotherapy (≤Grade 1). Monitor complete blood count for cytopenia at baseline and monthly thereafter for clinically significant changes during treatment. For prolonged hematological toxicities, interrupt LYNPARZA and monitor blood counts weekly until recovery. If the levels have not recovered to Grade 1 or less after 4 weeks, refer the patient to a hematologist for further investigations, including bone marrow analysis and blood sample for cytogenetics. Discontinue LYNPARZA if MDS/AML is confirmed.

IMPORTANT SAFETY INFORMATION (Cont'd)

WARNINGS AND PRECAUTIONS (Cont'd)

Pneumonitis: Including severe and fatal cases, has occurred in patients treated with LYNPARZA. In clinical studies, among patients who received LYNPARZA as a single agent or as part of a combination regimen, the incidence of pneumonitis, including fatal cases, was 1.0% (29/2851). If patients present with new or worsening respiratory symptoms such as dyspnea, cough, and fever, or a radiological abnormality occurs, interrupt LYNPARZA treatment and promptly assess the source of the symptoms. If pneumonitis is confirmed, discontinue LYNPARZA treatment and treat the patient appropriately.

Venous Thromboembolism (VTE): Including severe or fatal pulmonary embolism (PE), occurred in patients treated with LYNPARZA. Monitor patients for signs and symptoms of venous thrombosis and pulmonary embolism, and treat as medically appropriate, which may include long-term anticoagulation as clinically indicated.

Hepatotoxicity, including Drug-Induced Liver Injury (DILI): Hepatotoxicity, including severe and potentially fatal cases of DILI has occurred in patients treated with LYNPARZA. Evaluate bilirubin and transaminases at baseline and throughout treatment with LYNPARZA. For patients who develop abnormal liver tests after LYNPARZA, monitor more frequently for liver test abnormalities and clinical signs and symptoms of hepatic toxicity. If DILI is suspected, withhold LYNPARZA. Upon confirmation of DILI, discontinue LYNPARZA.

Embryo-Fetal Toxicity: Based on its mechanism of action and findings in animals, LYNPARZA can cause fetal harm. Verify pregnancy status in females of reproductive potential prior to initiating treatment.

Females

Advise females of reproductive potential of the potential risk to a fetus and to use effective contraception during treatment and for 6 months following the last dose.

ADVERSE REACTIONS—First-Line Maintenance BRCaM Advanced Ovarian Cancer

Most common adverse reactions (all Grades) in $\geq 10\%$ of patients who received LYNPARZA in the **first-line maintenance setting** for **SOLO-1** were: nausea (77%), fatigue (67%), abdominal pain (45%), vomiting (40%), anemia (38%), diarrhea (37%), constipation (28%), upper respiratory tract infection/influenza/nasopharyngitis/bronchitis (28%), dysgeusia (26%), decreased appetite (20%), dizziness (20%), neutropenia (17%), dyspepsia (17%), dyspnea (15%), leukopenia (13%), urinary tract infection (13%), thrombocytopenia (11%), and stomatitis (11%).

Most common laboratory abnormalities (Grades 1-4) in $\geq 25\%$ of patients who received LYNPARZA in the **first-line maintenance setting** for **SOLO-1** were: decrease in hemoglobin (87%), increase in mean corpuscular volume (87%), decrease in leukocytes (70%), decrease in lymphocytes (67%), decrease in absolute neutrophil count (51%), decrease in platelets (35%), and increase in serum creatinine (34%).

ADVERSE REACTIONS—First-Line Maintenance Advanced Ovarian Cancer in Combination with Bevacizumab

Most common adverse reactions (Grades 1-4) in $\geq 10\%$ of patients treated with LYNPARZA/bevacizumab and at a $\geq 5\%$ frequency compared to placebo/bevacizumab in the **first-line maintenance setting** for **PAOLA-1** were: nausea (53%), fatigue (including asthenia) (53%), anemia (41%), lymphopenia (24%), vomiting (22%), and leukopenia (18%). In addition, the most common adverse reactions ($\geq 10\%$) for patients receiving LYNPARZA/bevacizumab irrespective of the frequency compared with the placebo/bevacizumab arm were: diarrhea (18%), neutropenia (18%), urinary tract infection (15%), and headache (14%).

In addition, venous thromboembolism occurred more commonly in patients receiving LYNPARZA/bevacizumab (5%) than in those receiving placebo/bevacizumab (1.9%).

Most common laboratory abnormalities (Grades 1-4) in $\geq 25\%$ of patients for LYNPARZA in combination with bevacizumab in the **first-line maintenance setting** for **PAOLA-1** were: decrease in hemoglobin (79%), decrease in lymphocytes (63%), increase in serum creatinine (61%), decrease in leukocytes (59%), decrease in absolute neutrophil count (35%), and decrease in platelets (35%).

ADVERSE REACTIONS—Maintenance gBRCAm Recurrent Ovarian Cancer

Most common adverse reactions (Grades 1-4) in $\geq 20\%$ of patients who received LYNPARZA in the **maintenance setting** for **SOLO-2** were: nausea (76%), fatigue (including asthenia) (66%), anemia (44%), vomiting (37%), nasopharyngitis/upper respiratory tract infection (URI)/sinusitis/rhinitis/influenza (36%), diarrhea (33%), arthralgia/myalgia (30%), dysgeusia (27%), headache (26%), decreased appetite (22%), and stomatitis (20%).

Most common laboratory abnormalities (Grades 1-4) in $\geq 25\%$ of patients who received LYNPARZA in the **maintenance setting** for **SOLO-2** were: increase in mean corpuscular volume (89%), decrease in hemoglobin (83%), decrease in leukocytes (69%), decrease in lymphocytes (67%), decrease in absolute neutrophil count (51%), increase in serum creatinine (44%), and decrease in platelets (42%).

DRUG INTERACTIONS

Anticancer Agents: Clinical studies of LYNPARZA with other myelosuppressive anticancer agents, including DNA-damaging agents, indicate a potentiation and prolongation of myelosuppressive toxicity.

CYP3A Inhibitors: Avoid coadministration of strong or moderate CYP3A inhibitors when using LYNPARZA. If a strong or moderate CYP3A inhibitor must be coadministered, reduce the dose of LYNPARZA. Advise patients to avoid grapefruit, grapefruit juice, Seville oranges, and Seville orange juice during LYNPARZA treatment.

CYP3A Inducers: Avoid coadministration of strong or moderate CYP3A inducers when using LYNPARZA.

USE IN SPECIFIC POPULATIONS

Lactation: No data are available regarding the presence of olaparib in human milk, its effects on the breastfed infant or on milk production. Because of the potential for serious adverse reactions in the breastfed infant, advise a lactating woman not to breastfeed during treatment with LYNPARZA and for 1 month after receiving the final dose.

Pediatric Use: The safety and efficacy of LYNPARZA have not been established in pediatric patients.

Hepatic Impairment: No adjustment to the starting dose is required in patients with mild or moderate hepatic impairment (Child-Pugh classification A and B). There are no data in patients with severe hepatic impairment (Child-Pugh classification C).

Renal Impairment: No dosage modification is recommended in patients with mild renal impairment (CLcr 51-80 mL/min estimated by Cockcroft-Gault). In patients with moderate renal impairment (CLcr 31-50 mL/min), reduce the dose of LYNPARZA to 200 mg twice daily. There are no data in patients with severe renal impairment or end-stage renal disease (CLcr ≤ 30 mL/min).

Please see complete Prescribing Information, including Medication Guide.

You may report side effects related to AstraZeneca products [here](#).

References 1. LYNPARZA® (olaparib) [prescribing information]. Wilmington, DE: AstraZeneca Pharmaceuticals LP; 2025. 2. Zejula® (niraparib) [prescribing information]. Research Triangle Park, NC: GlaxoSmithKline; 2025. 3. Rubraca® (rucaparib) [prescribing information]. Vienna, Austria: zr pharma& GmbH; 2024.

This product information is intended for US healthcare professionals only. LYNPARZA is a registered trademark of the AstraZeneca group of companies. ZEJULA is a registered trademark of GlaxoSmithKline. RUBRACA is a registered trademark of zr pharma& GmbH. ©2025 AstraZeneca. All rights reserved. US-102306 Last Updated 8/25